



The Administration of Paracox™

THIS POSTER IS INTENDED AS A GUIDE FOR THE ADMINISTRATION OF PARACOX. Detailed instructions are to be found in the user instruction leaflet supplied with the pack.

1

- DRINKING WATER SHOULD BE WITHDRAWN 1-2 HOURS BEFORE VACCINATION BY RAISING DRINKERS
- VACCINATE HEALTHY BIRDS ONLY

2

- MIX VACCINE WELL BY SHAKING PACK VIGOROUSLY
- CALCULATE VACCINE

$$\frac{\text{TOTAL CHICKENS IN PEN}}{\text{TOTAL DRINKERS IN PEN}} \times 0.1 = \text{amount of vaccine per drinker (ml)}$$

 _____ x 0.1 = _____ ml

3

- DO NOT ADMINISTER VACCINE INTO HEADER TANK
- ADMINISTER IN BELL DRINKERS

4

- ENSURE EVEN DISTRIBUTION OF VACCINE AROUND DRINKER

5

~~FEED WITH ANTI-COCCIDIALS~~

- BIRDS MUST NOT RECEIVE FEED CONTAINING DRUGS WITH ANTICOCCIDIAL ACTIVITY

6

- CONTACT WITH LITTER IS REQUIRED FOR EFFECTIVE VACCINATION
- DO NOT MOVE FLOCKS FROM HOUSE TO HOUSE UNTIL AT LEAST 5 WEEKS AFTER VACCINATION

NOTES